

Package ‘faraway’

August 29, 2013

Version 1.0.5

Date 2011-04-05

Author Julian Faraway <jjf23@bath.ac.uk>

Maintainer Julian Faraway <jjf23@bath.ac.uk>

Title Functions and datasets for books by Julian Faraway.

Description Books are ‘‘Practical Regression and ANOVA in R’’ on CRAN, ‘‘Linear Models with R’’ published in August 2004 by CRC press and ‘‘Extending the Linear Model with R’’ published by CRC press in December 2005.

Depends R (>= 2.10)

License GPL

URL <http://www.maths.bath.ac.uk/~jjf23/>

LazyData yes

Repository CRAN

Date/Publication 2011-04-05 14:47:05

NeedsCompilation no

R topics documented:

aatemp	4
abrasion	5
aflatoxin	5
africa	6
alfalfa	7
amlxray	7
babyfood	8
beetle	9
bliss	9

breaking	10
broccoli	10
cathedral	11
chicago	12
chiczip	12
chmiss	13
choccake	13
chredlin	14
clot	15
cmob	15
cns	16
coagulation	17
composite	17
cornnit	18
corrosion	18
cpd	19
Cpplot	19
ctsib	20
death	20
debt	21
diabetes	22
dicentric	23
divusa	24
drugpsy	24
dvisits	25
eco	26
eggprod	27
eggs	27
epilepsy	28
esdcomp	28
exa	29
exb	29
eyegrade	30
fat	30
femsmoke	31
fpe	32
fruitfly	33
gala	33
gavote	34
haireye	35
halfnorm	35
happy	36
hormone	37
hprice	37
hsb	38
ilogit	39
infmort	39
irrigation	40

jsp	41
kanga	41
lawn	42
leafblotch	42
logit	43
mammalsleep	44
maxadjr	45
meatspec	45
melanoma	46
motorins	47
neighbor	48
nels88	49
nepali	49
nes96	50
oatvar	51
odor	52
ohio	52
orings	53
ozone	53
parstum	54
peanut	55
penicillin	55
pima	56
pipeline	57
pneumo	57
potuse	58
prostate	59
prplot	60
psid	60
pulp	61
pvc	62
qqnorml	62
rabbit	63
ratdrink	64
rats	64
resceram	65
salmonella	65
sat	66
savings	67
seatpos	67
semicond	68
sexab	69
sexfun	69
solder	70
sono	71
soybean	71
spector	72
speedo	73

star	74
stat500	74
strongx	75
suicide	75
teengamb	76
toenail	76
troutegg	77
truck	78
turtle	79
twins	79
uncviet	80
uswages	81
vif	81
vision	82
wafer	83
wavesolder	83
wbca	84
weldstrength	85
wheat	86
Index	87

aatemp

Annual mean temperatures in Ann Arbor, Michigan

Description

The data comes from the U.S. Historical Climatology Network.

Usage

```
data(aatemp)
```

Format

A data frame with 115 observations on the following 2 variables.

year year from 1854 to 2000

temp annual mean temperatures in degrees F in Ann Arbor

Source

United States Historical Climatology Network: <http://www.ncdc.noaa.gov/oa/climate/research/ushcn/ushcn.html>

abrasion

Wear on materials according to type, run and position

Description

The composite data frame has 16 rows and 4 columns. Four materials were fed into a wear testing machine and the amount of wear recorded. Four samples could be processed at the same time and the position of these samples may be important. A Latin square design was used.

Usage

```
data(abrasion)
```

Format

This data frame contains the following columns:

run The run number 1-4

position The position number 1-4

material The material A-D

wear The wear measured loss of weight in 0.1mm over testing period

Source

The Design and Analysis of Industrial Experiments by O. Davies, 1954, published by Wiley

aflatoxin

aflatoxin dosage and liver cancer in lab animals

Description

Aflatoxin B1 was fed to lab animals at vary doses and the number responding with liver cancer recorded.

Usage

```
data(aflatoxin)
```

Format

A data frame with 6 observations on the following 3 variables.

dose dose in ppb

total number of test animals

tumor number with liver cancer

Source

Gaylor DW (1987) "Linear nonparametric upper limits for low dose extrapolation" ASA Proceedings of the Biopharmaceutical Section.

Examples

```
data(aflatoxin)
```

```
africa
```

```
military coups and politics in sub-Saharan Africa
```

Description

Data is a subset of a larger study on factors affecting regime stability in Sub-Saharan Africa

Usage

```
data(africa)
```

Format

A data frame with 47 observations on the following 9 variables.

miltcoup number of successful military coups from independence to 1989

oligarchy number years country ruled by military oligarchy from independence to 1989

pollib Political liberalization - 0 = no civil rights for political expression, 1 = limited civil rights for expression but right to form political parties, 2 = full civil rights

parties Number of legal political parties in 1993

pctvote Percent voting in last election

popn Population in millions in 1989

size Area in 1000 square km

numelec Total number of legislative and presidential elections

numregim Number of regime types

Source

Bratton, Michael, and Nicholas Van De Walle. 1997. "Political Regimes and Regime Transitions in Africa, 1910-1994." *Study Number 106996*. Ann Arbor: Inter-University Consortium for Political and Social Research.

References

"Bayesian Methods: A Social and Behavioral Sciences Approach" by Jeff Gill 2002.

alfalfa

Effects of seed inoculum, irrigation and shade on alfalfa yield

Description

The alfalfa data frame has 25 rows and 4 columns. Data comes from an experiment to test the effects of seed inoculum, irrigation and shade on alfalfa yield. A latin square design has been used.

Usage

```
data(alfalfa)
```

Format

This data frame contains the following columns:

shade Distance of location from tree line divided into 5 shade areas

irrigation Irrigation effect divided into 5 levels

inoculum Four types of seed inoculum, A-D with E as control.

yield Dry matter yield of alfalfa

Source

Petersen, R.G. 1994. Agricultural Field Experiments, Design and Analysis. Marcel Dekker, Inc., New York. Pages 70-74. 1994

amlxray

Match pair study for AML and Xray link

Description

A matched case control study carried out to investigate the connection between X-ray usage and acute myeloid leukemia in childhood. The pairs are matched by age, race and county of residence.

Usage

```
data(amlxray)
```

Format

A data frame with 238 observations on the following 11 variables.

ID a factor denoting the matched pairs

disease 0=control, 1=case

Sex F or M

downs Presence of Downs syndrome: no or yes

age Age in years

Mray Did the mother ever have an Xray: no or yes

MupRay Did the mother have an Xray of the upper body during pregnancy: no or yes

MlowRay Did the mother have an Xray of the lower body during pregnancy: no or yes

Fray Did the father ever have an Xray: no or yes

Cray Did the child ever have an Xray: no or yes

CnRay Total number of Xrays of the child 1=none < 2=1 or 2 < 3=3 or 4 < 4= 5 or more

Source

Chap T. Le (1998) "Applied Categorical Data Analysis" Wiley.

babyfood

Respiratory disease rates of babies fed in different ways

Description

Example Dataset from "Practical Regression and Anova"

Usage

```
data(babyfood)
```

Format

See for yourself

Source

See Reference

References

Reference details may be found in "Practical Regression and Anova" by Julian Faraway

beetle	<i>Beetles exposed to fumigant</i>
--------	------------------------------------

Description

Example Dataset from "Practical Regression and Anova"

Usage

```
data(beetle)
```

Format

See for yourself

Source

See Reference

References

Reference details may be found in "Practical Regression and Anova" by Julian Faraway

bliss	<i>Bliss insecticide data</i>
-------	-------------------------------

Description

Example Dataset from "Practical Regression and Anova"

Usage

```
data(bliss)
```

Format

See for yourself

Source

See Reference

References

Reference details may be found in "Practical Regression and Anova" by Julian Faraway

 breaking

Breaking strength of materials

Description

An experiment was conducted to select the supplier of raw materials for production of a component. The breaking strength of the component was the objective of interest. Four suppliers were considered. The four operators can only produce one component each per day. A Latin square design was used.

Usage

```
data(breaking)
```

Format

A data frame with 16 observations on the following 4 variables.

y The breaking strength of the component

operator the operator - a factor with levels op1 op2 op3 op4

day the day of production - a factor with levels day1 day2 day3 day4

supplier the supplier of the raw material - a factor with levels A B C D

Source

Lentner M. and Bishop T. (1986) Experimental Design and Analysis, Valley Book Company

 broccoli

Broccoli weight variation

Description

A number of growers supply broccoli to a food processing plant. The plant instructs the growers to pack the broccoli into standard size boxes. There should be 18 clusters of broccoli per box and each cluster should weigh between 1.33 and 1.5 pounds. Because the growers use different varieties, methods of cultivation etc, there is some variation in the cluster weights. The plant manager selected 3 growers at random and then 4 boxes at random supplied by these growers. 3 clusters were selected from each box.

Usage

```
data(broccoli)
```

Format

A data frame with 36 observations on the following 4 variables.

wt weight of broccoli

grower the grower - a factor with levels 1 2 3

box the box - a factor with levels 1 2 3 4

cluster the cluster - a factor with levels 1 2 3

Source

Lentner M. and Bishop T. (1986) Experimental Design and Analysis, Valley Book Company

cathedral	<i>Cathedral nave heights and lengths in England</i>
-----------	--

Description

Example Dataset from "Practical Regression and Anova"

Usage

```
data(cathedral)
```

Format

See for yourself

Source

See Reference

References

Reference details may be found in "Practical Regression and Anova" by Julian Faraway

chicago

Chicago insurance redlining

Description

Example Dataset from "Practical Regression and Anova"

Usage

```
data(chicago)
```

Format

See for yourself

Source

See Reference

References

Reference details may be found in "Practical Regression and Anova" by Julian Faraway

chiczip

Chicago zip codes north-south

Description

Example Dataset from "Practical Regression and Anova"

Usage

```
data(chiczip)
```

Format

See for yourself

Source

See Reference

References

Reference details may be found in "Practical Regression and Anova" by Julian Faraway

chmiss	<i>Chicago insurance redlining</i>
--------	------------------------------------

Description

Data from a 1970's study on the relationship between insurance redlining in Chicago and racial composition, fire and theft rates, age of housing and income in 47 zip codes. Missing values have been randomly added.

Usage

```
data(chmiss)
```

Format

This dataframe contains the following columns

race racial composition in percent minority

fire fires per 100 housing units

theft theft per 1000 population

age percent of housing units built before 1939

involact new FAIR plan policies and renewals per 100 housing units

income median family income in thousands of dollars

side North or South side of Chicago

Source

Adapted from "Data : A Collection of Problems from Many Fields for the Student and Research Worker" by D. Andrews and A. Herzberg published by Springer-Verlag, in 1985

choccake	<i>Chocolate cake experiment with split plot design</i>
----------	---

Description

An experiment was conducted to determine the effect of recipe and baking temperature on chocolate cake quality. 15 batches of cake mix for each recipe were prepared. Each batch was sufficient for six cakes. Each of the six cakes was baked at a different temperature which was randomly assigned. Several measures of cake quality were recorded of which breaking angle was just one.

Usage

```
data(choccake)
```

Format

A data frame with 270 observations on the following 4 variables.

recipe Chocolate for recipe 1 was added at 40C, Chocolate for recipe 2 was added at 60C and recipe 3 had extra sugar

batch batch number from 1 to 15

temp temperature at which cake was baked: 175C 185C 195C 205C 215C 225C

breakang the breaking angle of the cake

Source

Cochran W. and Cox G. (1992) Experimental Designs, 2nd Edition Wiley

chredlin

Chicago insurance redlining

Description

Data from a 1970's study on the relationship between insurance redlining in Chicago and racial composition, fire and theft rates, age of housing and income in 47 zip codes

Usage

```
data(chredlin)
```

Format

This dataframe contains the following columns

race racial composition in percent minority

fire fires per 100 housing units

theft theft per 1000 population

age percent of housing units built before 1939

involact new FAIR plan policies and renewals per 100 housing units

income median family income in thousands of dollars

side North or South side of Chicago

Source

Adapted from "Data : A Collection of Problems from Many Fields for the Student and Research Worker" by D. Andrews and A. Herzberg published by Springer-Verlag, in 1985

clot

Blood clotting times

Description

Example Dataset from "Practical Regression and Anova"

Usage

```
data(clot)
```

Format

See for yourself

Source

See Reference

References

Reference details may be found in "Practical Regression and Anova" by Julian Faraway

cmob

Social class mobility from 1971 to 1981 in the UK

Description

Social class mobility from 1971 to 1981 for 42425 men from the United Kingdom census. Subjects were aged 45-64.

Usage

```
data(cmob)
```

Format

A data frame with 36 observations on the following 3 variables.

y Frequency of observation

class71 social class in 1971 - a factor with levels I, professionals, II semi-professionals, IIIN skilled non-manual, IIIM skilled manual, IV semi-skilled, V unskilled

class81 social class in 1981 - a factor with levels I II IIIN IIIM IV V with same classification

Source

D. Blane and S. Harding and M. Rosato (1999) "Does social mobility affect the size of the socioeconomic mortality differential?: Evidence from the Office for National Statistics Longitudinal Study" JRSS-A, 162 59-70.

 cns

Malformations of the central nervous system

Description

Frequencies of various malformations of the central nervous system recorded on live births in South Wales, UK. Study was designed to determine the effect of water hardness on the incidence of such malformations.

Usage

```
data(cns)
```

Format

A data frame with 16 observations on the following 7 variables.

Area a factor with levels Cardiff GlamorganC GlamorganE GlamorganW MonmouthOther MonmouthV Newport Swansea being areas of South Wales

NoCNS count of births with no CNS problem

An count of Anencephalus births

Sp count of Spina Bifida births

Other count of other CNS births

Water water hardness

Work a factor with levels Manual NonManual being the type of work done by the parents

Source

C. Lowe and C. Roberts and S. Lloyd, (1971) Malformations of the central nervous system and softness of local water supplies, British Medical Journal, 15,357-361.

References

P. McCullagh and J. Nelder (1989), Generalized Linear Models, Chapman and Hall, 2nd Ed.

coagulation	<i>Blood coagulation times by diet</i>
-------------	--

Description

Dataset comes from a study of blood coagulation times. 24 animals were randomly assigned to four different diets and the samples were taken in a random order.

Usage

```
data(coagulation)
```

Format

This dataframe contains the following columns

```
coag  coagulation time in seconds  
diet  diet type - A,B,C or D
```

Source

"Statistics for Experimenters" by G. P. Box, W. G. Hunter and J. S. Hunter, Wiley, 1978

composite	<i>Strength of a thermoplastic composite depending on two factors</i>
-----------	---

Description

The composite data frame has 9 rows and 3 columns. Data comes from an experiment to test the strength of a thermoplastic composite depending on the power of a laser and speed of a tape.

Usage

```
data(composite)
```

Format

This data frame contains the following columns:

```
strength  interply bond strength of the composite  
laser     laser power at 40, 50 or 60W  
tape     tape speed, slow=6.42 m/s, medium=13m/s and fast=27m/s
```

Source

Mazumdar, S and Hoa S (1995) "Application of a Taguchi Method for Process enhancement of an online consolidation technique" Composites 26, 669-673

 cornnit

Corn yields from nitrogen application

Description

The relationship between corn yield (bushels per acre) and nitrogen (pounds per acre) fertilizer application were studied in Wisconsin.

Usage

```
data(cornnit)
```

Format

A data frame with 44 observations on the following 2 variables.

yield corn yield in bushels per acre

nitrogen pounds per acre

Source

Unknown

 corrosion

Corrosion loss in Cu-Ni alloys

Description

Data consist of thirteen specimens of 90/10 Cu-Ni alloys with varying iron content in percent. The specimens were submerged in sea water for 60 days and the weight loss due to corrosion was recorded in units of milligrams per square decimeter per day.

Usage

```
data(corrosion)
```

Format

This dataframe contains the following columns

Fe Iron content in percent

loss Weight loss in mg per square decimeter per day

Source

"Applied Regression Analysis" by N. Draper and H. Smith, Wiley, 1998

`cpd`*Projected and actual sales of 20 consumer products*

Description

Projected and actual sales of 20 consumer products. Data have been disguised from original form.

Usage

```
data(cpd)
```

Format

A data frame with 20 observations on the following 2 variables.

projected projected sales in dollars

actual actual sales in dollars

Source

G. Whitmore (1986) "Inverse Gaussian Ratio Estimation" Applied Statistics, 35, 8-15.

`Cpplot`*Cp plot*

Description

Makes a Cp plot

Usage

```
Cpplot(cp)
```

Arguments

`cp` A leaps object returned from `leaps()`

Details

Requires leaps package

Value

none

Author(s)

Julian Faraway

See Also

leaps()

ctsib

Effects of surface and vision on balance

Description

Example Dataset from "Practical Regression and Anova"

Usage

data(ctsib)

Format

See for yourself

Source

See Reference

References

Reference details may be found in "Practical Regression and Anova" by Julian Faraway

death

Death penalty in Florida 1977

Description

Data on 326 defendants in homicide indictments in 20 Florida counties during 1976-77.

Usage

data(death)

Format

A data frame with 8 observations on the following 4 variables.

`y` a numeric vector

`penalty` Did the subject receive the death penalty? no or yes

`victim` Was the victim black or white?

`defend` Was the defendant black or white?

Source

Radelet M. (1981) Racial characteristics and the imposition of the death penalty. *Amer. Sociol. Rev.* **46** 918-927.

References

Agresti A. (1990) *Categorical Data Analysis*, Wiley.

 debt

psychology of debt

Description

The data arise from a large postal survey on the psychology of debt.

Usage

`data(debt)`

Format

A data frame with 464 observations on the following 13 variables.

incomegp income group (1=lowest, 5=highest)

house security of housing tenure (1=rent, 2=mortgage, 3=owned outright)

children number of children in household

singpar is the respondent a single parent?

agegp age group (1=youngest)

bankacc does the respondent have a bank account?

bsocacc does the respondent have a building society account?

manage self-rating of money management skill (high values=high skill)

ccarduse how often did s/he use credit cards (1=never... 3=regularly)

cigbuy does s/he buy cigarettes?

xmasbuy does s/he buy Christmas presents for children?

locintrn score on a locus of control scale (high values=internal)

prodebt score on a scale of attitudes to debt (high values=favourable to debt)

Details

All yes/no questions are coded 0=no, 1=yes. Locus of control is a personality measure introduced by Rotter, which claims to differentiate people according to how much they feel things that happen to them are as a result of processes within themselves (internal locus of control) or outside events (external locus of control).

Source

Lea, Webley & Walker, 1995, Journal of Economic Psychology, 16, 181-201 Data obtained from <http://www.exeter.ac.uk/~SEGLea>.

diabetes

Diabetes and obesity, cardiovascular risk factors

Description

403 African Americans were interviewed in a study to understand the prevalence of obesity, diabetes, and other cardiovascular risk factors in central Virginia.

Usage

```
data(diabetes)
```

Format

A data frame with 403 observations on the following 19 variables.

```
id Subject ID
chol Total Cholesterol
stab.glu Stabilized Glucose
hdl High Density Lipoprotein
ratio Cholesterol/HDL Ratio
glyhb Glycosolated Hemoglobin
location County - a factor with levels Buckingham Louisa
age age in years
gender a factor with levels male female
height height in inches
weight weight in pounds
frame a factor with levels small medium large
bp.1s First Systolic Blood Pressure
bp.1d First Diastolic Blood Pressure
bp.2s Second Systolic Blood Pressure
bp.2d Second Diastolic Blood Pressure
waist waist in inches
hip hip in inches
time.ppn Postprandial Time (in minutes) when Labs were Drawn
```

Details

Glycosolated hemoglobin greater than 7.0 is usually taken as a positive diagnosis of diabetes

Source

Willems JP, Saunders JT, DE Hunt, JB Schorling: Prevalence of coronary heart disease risk factors among rural blacks: A community-based study. *Southern Medical Journal* 90:814-820; 1997

References

Schorling JB, Roach J, Siegel M, Baturka N, Hunt DE, Guterbock TM, Stewart HL: A trial of church-based smoking cessation interventions for rural African Americans. *Preventive Medicine* 26:92-101; 1997

 dicentric

Radiation dose effects on chromosomal abnormality

Description

An experiment was conducted to determine the effect of gamma radiation on the numbers of chromosomal abnormalities observed

Usage

data(dicentric)

Format

A data frame with 27 observations on the following 4 variables.

cells Number of cells in hundreds

ca Number of chromosomal abnormalities

doseamt amount of dose in Grays

doserate rate of dose in Grays/hour

Source

Puott R. and Reeder E. (1976) The effect of changes in dose rate on the yield of chromosome aberrations in human lymphocytes exposed to gamma radiation. *Mutation Research*. 35, 437-444.

References

Frome E. and DuFrain R. (1986) Maximum Likelihood Estimation for Cytogenic Dose-Response Curves. *Biometrics*. 42, 73-84.

divusa	<i>Divorce in the USA 1920-1996</i>
--------	-------------------------------------

Description

Divorce rates in the USA from 1920-1996

Usage

```
data(divusa)
```

Format

A data frame with 77 observations on the following 7 variables.

year the year from 1920-1996

divorce divorce per 1000 women aged 15 or more

unemployed unemployment rate

femlab percent female participation in labor force aged 16+

marriage marriages per 1000 unmarried women aged 16+

birth births per 1000 women aged 15-44

military military personnel per 1000 population

Source

Unknown

drugpsy	<i>Choice of drug treatment for psychiatry patients</i>
---------	---

Description

A sample of psychiatry patients were cross-classified by their diagnosis and whether a drug treatment was prescribed.

Usage

```
data(drugpsy)
```

Format

A data frame with 10 observations on the following 3 variables.

y the number of patients

diagnosis a factor with levels Affective.Disorder Neurosis Personality.Disorder Schizophrenia Special.Symptoms

drug a factor with levels no yes

Source

Helmes E. and Fekken G. (1986) Effects of psychotropic drugs and psychiatric illness on vocational aptitude and interest assessment. *J. Clin. Psychol.* **42** 569-576

References

Agresti A. (1990) "Categorical Data Analysis" Wiley

dvisits

Doctor visits in Australia

Description

The data come from the Australian Health Survey of 1977-78 and consist of 5190 single adults where young and old have been oversampled.

Usage

```
data(dvisits)
```

Format

A data frame with 5190 observations on the following 19 variables.

sex 1 if female, 0 if male

age Age in years divided by 100 (measured as mid-point of 10 age groups from 15-19 years to 65-69 with 70 or more coded treated as 72)

agesq age squared

income Annual income in Australian dollars divided by 1000 (measured as mid-point of coded ranges Nil, less than 200, 200-1000, 1001-, 2001-, 3001-, 4001-, 5001-, 6001-, 7001-, 8001-10000, 10001-12000, 12001-14000, with 14001- treated as 15000)

levyplus 1 if covered by private health insurance fund for private patient in public hospital (with doctor of choice), 0 otherwise

freepoor 1 if covered by government because low income, recent immigrant, unemployed, 0 otherwise

freerepa 1 if covered free by government because of old-age or disability pension, or because invalid veteran or family of deceased veteran, 0 otherwise

illness Number of illnesses in past 2 weeks with 5 or more coded as 5

actdays Number of days of reduced activity in past two weeks due to illness or injury

hscore General health questionnaire score using Goldberg's method. High score indicates bad health

chcond1 1 if chronic condition(s) but not limited in activity, 0 otherwise

chcond2 1 if chronic condition(s) and limited in activity, 0 otherwise

doctorco Number of consultations with a doctor or specialist in the past 2 weeks

nondocco Number of consultations with non-doctor health professionals (chemist, optician, physiotherapist, social worker, district community nurse, chiropodist or chiropractor) in the past 2 weeks

hospadmi Number of admissions to a hospital, psychiatric hospital, nursing or convalescent home in the past 12 months (up to 5 or more admissions which is coded as 5)

hospdays Number of nights in a hospital, etc. during most recent admission: taken, where appropriate, as the mid-point of the intervals 1, 2, 3, 4, 5, 6, 7, 8-14, 15-30, 31-60, 61-79 with 80 or more admissions coded as 80. If no admission in past 12 months then equals zero

medicine Total number of prescribed and nonprescribed medications used in past 2 days

prescrib Total number of prescribed medications used in past 2 days

nonpresc Total number of nonprescribed medications used in past 2 days

Source

Cameron A, Trivedi P, Milne F and Piggot J (1988) A Microeconomic model of the demand for health care and health insurance in Australia, *Review of Economic Studies* 55, 85-106

eco

Ecological regression example

Description

Relationship between 1998 per capita income dollars from all sources and the proportion of legal state residents born in the United States in 1990 for each of the 50 states plus the District of Columbia

Usage

`data(eco)`

Format

This dataframe contains the following columns

usborn Percentage of population born in the United States

income Per capita annual income in dollars

home Percentage born in state

pop Population of state

Source

US Bureau of the Census

eggprod	<i>Treatment and block effects on egg production</i>
---------	--

Description

The composite data frame has 12 rows and 3 columns. Six pullets were placed into each of 12 pens. Four blocks were formed from groups of 3 pens based on location. Three treatments were applied. The number of eggs produced was recorded

Usage

```
data(eggprod)
```

Format

This data frame contains the following columns:

treat Three treatments: O, E or F

block Four blocks labeled 1-4

eggs Number of eggs produced

Source

Mead, R., R.N. Curnow, and A.M. Hasted. 1993. *Statistical Methods in Agriculture and Experimental Biology*. Chapman and Hall, London, p. 64. 1993

eggs	<i>Laboratory testing of dried egg fat content</i>
------	--

Description

Example Dataset from "Practical Regression and Anova"

Usage

```
data(eggs)
```

Format

See for yourself

Source

See Reference

References

Reference details may be found in "Practical Regression and Anova" by Julian Faraway

epilepsy

Seizure rates of epileptics under treatment

Description

Example Dataset from "Practical Regression and Anova"

Usage

```
data(epilepsy)
```

Format

See for yourself

Source

See Reference

References

Reference details may be found in "Practical Regression and Anova" by Julian Faraway

esdcomp

Complaints about emergency room doctors

Description

Data was recorded on 44 doctors working in an emergency service at a hospital to study the factors affecting the number of complaints received.

Usage

```
data(esdcomp)
```

Format

A data frame with 44 observations on the following 6 variables.

visits the number of patient visits

complaints the number of complaints

residency is the doctor in residency training N or Y

gender gender of doctor F or M

revenue dollars per hour earned by the doctor

hours total number of hours worked

Source

Chap T. Le (1998) "Applied Categorical Data Analysis" Wiley

exa

Non parametric regression test data A

Description

Example Dataset from "Practical Regression and Anova"

Usage

```
data(exa)
```

Format

See for yourself

Source

See Reference

References

Reference details may be found in "Practical Regression and Anova" by Julian Faraway

exb

Non parametric regression test data B

Description

Example Dataset from "Practical Regression and Anova"

Usage

```
data(exb)
```

Format

See for yourself

Source

See Reference

References

Reference details may be found in "Practical Regression and Anova" by Julian Faraway

eyegrade	<i>grading of eye pairs for distance vision</i>
----------	---

Description

A sample of women are rated for the performance of distance vision in each eye.

Usage

data(eyegrade)

Format

A data frame with 16 observations on the following 3 variables.

y the observed count

right rated vision in the right eye - a factor with levels best second third worst

left rated vision in the left eye - a factor with levels best second third worst

Source

A. Stuart (1955) A test for homogeneity of the marginal distributions in a two-way classification, *Biometrika*, 42, 412-416.

fat	<i>Percentage of Body Fat and Body Measurements</i>
-----	---

Description

Age, weight, height, and 10 body circumference measurements are recorded for 252 men. Each man's percentage of body fat was accurately estimated by an underwater weighing technique.

Usage

data(fat)

Format

A data frame with 252 observations on the following 18 variables.

brozek Percent body fat using Brozek's equation, 457/Density - 414.2

siri Percent body fat using Siri's equation, 495/Density - 450

density Density (gm/cm³)

age Age (yrs)

weight Weight (lbs)

height Height (inches)
 adipos Adiposity index = $\text{Weight}/\text{Height}^2$ (kg/m²)
 free Fat Free Weight = (1 - fraction of body fat) * Weight, using Brozek's formula (lbs)
 neck Neck circumference (cm)
 chest Chest circumference (cm)
 abdom Abdomen circumference (cm) at the umbilicus and level with the iliac crest
 hip Hip circumference (cm)
 thigh Thigh circumference (cm)
 knee Knee circumference (cm)
 ankle Ankle circumference (cm)
 biceps Extended biceps circumference (cm)
 forearm Forearm circumference (cm)
 wrist Wrist circumference (cm) distal to the styloid processes

Source

Johnson R. Journal of Statistics Education v.4, n.1 (1996)

femsmoke

Mortality due to smoking according age group in women

Description

In 1972-74, a survey of one in six residents of Wickham, near Newcastle, England was made. Twenty years later, this data recorded in a follow-up study. Only women who are current smokers or who have never smoked are included.

Usage

data(femsmoke)

Format

A data frame with 28 observations on the following 4 variables.

y observed count for given combination

smoker a factor with levels yes no

dead a factor with levels yes no

age a factor with agegroup levels 18-24 25-34 35-44 45-54 55-64 65-74 75+

Source

D. Appleton, J. French, M. Vanderpump (1996) "Ignoring a Covariate: An Example of Simpson's Paradox" American Statistician, 50, 340-341

fpe

1981 French Presidential Election

Description

Elections for the French presidency proceed in two rounds. In 1981, there were 10 candidates in the first round. The top two candidates then went on to the second round, which was won by Francois Mitterand over Valery Giscard-d'Estaing. The losers in the first round can gain political favors by urging their supporters to vote for one of the two finalists. Since voting is private, we cannot know how these votes were transferred, we might hope to infer from the published vote totals how this might have happened. Data is given for vote totals in every fourth department of France:

Usage

```
data(fpe)
```

Format

This dataframe contains the following columns (vote totals are in thousands)

EI Electeur Inscrits (registered voters)

A Voters for Mitterand in the first round

B Voters for Giscard in the first round

C Voters for Chirac in the first round

D Voters for Communists in the first round

E Voters for Ecology party in the first round

F Voters for party F in the first round

G Voters for party G in the first round

H Voters for party H in the first round

I Voters for party I in the first round

J Voters for party J in the first round

K Voters for party K in the first round

A2 Voters for Mitterand in the second round

B2 Voters for party Giscard in the second round

N Difference between the number of voters in the second round and in the first round

Source

"The Teaching of Practical Statistics" by C.W. Anderson and R.M. Loynes, Wiley, 1987

fruitfly

Longevity of fruitflies depending on sexual activity and thorax length

Description

The fruitfly data frame has 9 rows and 3 columns. 125 fruitflies were divided randomly into 5 groups of 25 each. The response was the longevity of the fruitfly in days. One group was kept solitary, while another was kept individually with a virgin female each day. Another group was given 8 virgin females per day. As an additional control the fourth and fifth groups were kept with one or eight pregnant females per day. Pregnant fruitflies will not mate. The thorax length of each male was measured as this was known to affect longevity. One observation in the many group has been lost.

Usage

```
data(fruitfly)
```

Format

This data frame contains the following columns:

thorax Thorax length

longevity Lifetime in days

activity The group: isolated = fly kept solitary, one = fly kept with one pregnant fruitfly, many = fly kept with eight pregnant fruitflies, low = fly kept with one virgin fruitfly, high = fly kept with eight virgin fruitflies.

Source

"Sexual Activity and the Lifespan of Male Fruitflies" by L. Partridge and M. Farquhar, Nature, 1981, 580-581

gala

Species diversity on the Galapagos Islands

Description

There are 30 Galapagos islands and 7 variables in the dataset. The relationship between the number of plant species and several geographic variables is of interest. The original dataset contained several missing values which have been filled for convenience.

Usage

```
data(gala)
```

Format

The dataset contains the following variables

Species the number of plant species found on the island

Endemics the number of endemic species

Area the area of the island (km²)

Elevation the highest elevation of the island (m)

Nearest the distance from the nearest island (km)

Scruz the distance from Santa Cruz island (km)

Adjacent the area of the adjacent island (square km)

Source

M. P. Johnson and P. H. Raven (1973) "Species number and endemism: The Galapagos Archipelago revisited" *Science*, 179, 893-895

gavote

Undercounted votes in Georgia in 2000 presidential election

Description

The data comes from the US presidential election in the state of Georgia. The undercount is the difference between the number of ballots cast and votes recorded. Voters may have chosen not to vote for president, voted for more than one candidate (disqualified) or the equipment may have failed to register their choice.

Usage

data(gavote)

Format

A data frame with 159 observations on the following 10 variables. Each case represents a county in Georgia.

equip The voting equipment used: LEVER, OS-CC (optical, central count), OS-PC (optical, precinct count) PAPER, PUNCH

econ economic status of county: middle poor rich

perAA percent of African Americans in county

rural indicator of whether county is rural or urban

atlanta indicator of whether county is in Atlanta or not: notAtlanta

gore number of votes for Gore

bush number of votes for Bush

other number of votes for other candidates

votes number of votes

ballots number of ballots

Source

Meyer M. (2002) Uncounted Votes: Does Voting Equipment Matter? *Chance*, 15(4), 33-38

haireye	<i>Hair and eye color</i>
---------	---------------------------

Description

Data collected from 592 students in an introductory statistics class

Usage

```
data(haireye)
```

Format

A data frame with 16 observations on the following 3 variables.

y count of the number of student with given hair/eye combination

eye a factor with levels green hazel blue brown

hair a factor with levels BLACK BROWN RED BLOND

Source

Snee R. (1974) Graphical display of two-way contingency tables. *American Statistician*, 28, 9-12

halfnorm	<i>Half Normal Plot</i>
----------	-------------------------

Description

Makes a half-normal plot

Usage

```
halfnorm(x, nlab = 2, labs = as.character(1:length(x)), ylab = "Sorted Data",
...)
```

Arguments

x	a numeric vector
nlab	number of points to label
labs	labels for points
ylab	label for Y-axis
...	arguments passed to plot()

Value

none

Author(s)

Julian Faraway

See Also

qqnorm

Examples

```
halfnorm(runif(10))
```

happy

love, work and happiness

Description

Data were collected from 39 students in a University of Chicago MBA class

Usage

```
data(happy)
```

Format

A data frame with 39 observations on the following 5 variables.

happy Happiness on a 10 point scale where 10 is most happy

money family income in thousands of dollars

sex 1 = satisfactory sexual activity, 0 = not

love 1 = lonely, 2 = secure relationships, 3 = deep feeling of belonging and caring

work 5 point scale where 1 = no job, 3 = OK job, 5 = great job

Source

George and McCulloch (1993) "Variable Selection via Gibbs Sampling" JASA, 88, 881-889

hormone	<i>Hormones and Sexual Orientation</i>
---------	--

Description

Example Dataset from "Practical Regression and Anova"

Usage

```
data(hormone)
```

Format

See for yourself

Source

See Reference

References

Reference details may be found in "Practical Regression and Anova" by Julian Faraway

hprice	<i>Housing prices in US cities 86-94</i>
--------	--

Description

Data on housing prices in 36 US metropolitan statistical areas (MSAs) over 9 years from 1986-1994 were collected.

Usage

```
data(hprice)
```

Format

A data frame with 324 observations on the following 8 variables.

lvsale log average sale price

lvcap average per capita income

lvgrow percentage growth in per capita income

lvrtest Regulatory environment index (high values = more regulations)

lvrctrl Rent control - a factor with levels 0=no 1=yes

lvrcoast Adjacent to a coastline - a factor with levels 0=no 1=yes

lvrmsa indicator for the MSA

lvrtime Year 1=1986 to 9=1994

Source

Longitudinal and Panel Data: Analysis and Applications in the Social Sciences, by Edward W. Frees, Cambridge University Press, August 2004.

hsb

Career choice of high school students

Description

Data was collected as a subset of the "High School and Beyond" study conducted by the National Education Longitudinal Studies (NELS) program of the National Center for Education Statistics (NCES).

Usage

data(hsb)

Format

A data frame with 200 observations on the following 11 variables.

id ID of student

gender a factor with levels female male

race a factor with levels african-amer asian hispanic white

ses socioeconomic class - a factor with levels high low middle

schtyp school type - a factor with levels private public

prog choice of high school program - a factor with levels academic general vocation

read reading score

write writing score

math math score

science science score

socst social science score

Details

One purpose of the study was to determine which factors are related to the choice of the type of program, academic, vocational or general, that the students pursue in high school.

Source

National Education Longitudinal Studies (NELS) program of the National Center for Education Statistics (NCES).

ilomit	<i>Inverse Logit Transformation</i>
--------	-------------------------------------

Description

Computes the inverse logit transformation

Usage

```
ilomit(x)
```

Arguments

x a numeric vector

Value

$\exp(x)/(1+\exp(x))$

Author(s)

Julian Faraway

See Also

logit

Examples

```
ilomit(1:3)
#[1] 0.7310586 0.8807971 0.9525741
```

infmort	<i>Infant mortality according to income and region</i>
---------	--

Description

The infmort data frame has 105 rows and 4 columns. The infant mortality in regions of the world may be related to per capita income and whether oil is exported. The dataset is not recent.

Usage

```
data(infmort)
```

Format

This data frame contains the following columns:

region Region of the world, Africa, Europe, Asia or the Americas

income Per capita annual income in dollars

mortality Infant mortality in deaths per 1000 births

oil Does the country export oil or not?

Source

Unknown

irrigation	<i>Agricultural experiment with irrigation</i>
------------	--

Description

Example Dataset from "Practical Regression and Anova"

Usage

```
data(irrigation)
```

Format

See for yourself

Source

See Reference

References

Reference details may be found in "Practical Regression and Anova" by Julian Faraway

jsp

Junior Schools Project

Description

Example Dataset from "Practical Regression and Anova"

Usage

```
data(jsp)
```

Format

See for yourself

Source

See Reference

References

Reference details may be found in "Practical Regression and Anova" by Julian Faraway

kanga

Historic Kangaroos

Description

Example Dataset from "Practical Regression and Anova"

Usage

```
data(kanga)
```

Format

See for yourself

Source

See Reference

References

Reference details may be found in "Practical Regression and Anova" by Julian Faraway

lawn	<i>Cut-off times of lawnmowers</i>
------	------------------------------------

Description

Data on the cut-off times of lawnmowers was collected. 3 machines were randomly selected from those produced by manufacturers A and B. Each machine was tested twice at low speed and high speed.

Usage

```
data(lawn)
```

Format

A data frame with 24 observations on the following 4 variables.

```
manufact Manufacturer - a factor with levels A B
machine Lawn mower - a factor with levels m1 m2 m3 m4 m5 m6
speed Speed of testing - a factor with levels H L
time cut-off time
```

Source

Unknown.

leafblotch	<i>Leaf blotch on barley</i>
------------	------------------------------

Description

The data gives the proportion of leaf area affected by leaf blotch on 10 varieties of barley at 9 different sites.

Usage

```
data(leafblotch)
```

Format

A data frame with 90 observations on the following 3 variables.

```
blotch proportion of the barley leaf affected by blotch
site the physical location - a factor with levels 1 2 3 4 5 6 7 8 9
variety variety of barley - a factor with levels 1 2 3 4 5 6 7 8 9 10
```

Source

R. W. M. Wedderburn (1974) "Quasilikelihood functions, generalized linear models and the Gauss-Newton method" *Biometrika*, 61, 439-447.

References

P. McCullagh and J. Nelder (1989) "Generalized Linear Models" Chapman and Hall, 2nd ed.

logit	<i>Logit transformation</i>
-------	-----------------------------

Description

Computes the logit transformation

Usage

```
logit(x)
```

Arguments

x a numeric vector

Details

x <=0 or >=1 will return NA

Value

$\log(x/(1-x))$

Author(s)

Julian Faraway

See Also

ilogit

Examples

```
logit(c(0.1,0.5,1.0,1.1))
#[1] -2.197225  0.000000      NA      NA
```

mammalsleep

Sleep in Mammals: Ecological and Constitutional Correlates

Description

The mammalsleep data frame has 62 rows and 10 columns. Sleep in Mammals: Ecological and Constitutional Correlates

Usage

```
data(mammalsleep)
```

Format

This data frame contains the following columns:

body body weight in kg

brain brain weight in g

nondream slow wave ("nondreaming") sleep (hrs/day)

dream paradoxical ("dreaming") sleep (hrs/day)

sleep total sleep (hrs/day) (sum of slow wave and paradoxical sleep)

lifespan maximum life span (years)

gestation gestation time (days)

predation predation index (1-5) 1 = minimum (least likely to be preyed upon) to 5 = maximum (most likely to be preyed upon)

exposure sleep exposure index (1-5) 1 = least exposed (e.g. animal sleeps in a well-protected den) 5 = most exposed

danger overall danger index (1-5) (based on the above two indices and other information) 1 = least danger (from other animals) 5 = most danger (from other animals)

Source

"Sleep in Mammals: Ecological and Constitutional Correlates" by Allison, T. and Cicchetti, D. (1976), Science, November 12, vol. 194, pp. 732-734.

maxadjr	<i>Maximum Adjusted R-squared</i>
---------	-----------------------------------

Description

Displays the best models from a leaps object

Usage

```
maxadjr(1, best=3)
```

Arguments

1	A leaps object returned from leaps()
best	An optional argument specify the number of models to be returned taking the default value of 3

Details

Requires leaps package

Value

A list of the best models

Author(s)

Julian Faraway

See Also

leaps()

meatspec	<i>Meat spectrometry to determine fat content</i>
----------	---

Description

A Tecator Infratec Food and Feed Analyzer working in the wavelength range 850 - 1050 nm by the Near Infrared Transmission (NIT) principle was used to collect data on samples of finely chopped pure meat. 215 samples were measured. For each sample, the fat content was measured along with a 100 channel spectrum of absorbances. Since determining the fat content via analytical chemistry is time consuming we would like to build a model to predict the fat content of new samples using the 100 absorbances which can be measured more easily.

Usage

```
data(meatspec)
```

Format

Dataset contains the following variables

V1-V100 absorbances across a range of 100 wavelengths

fat fat content

Source

H. H. Thodberg (1993) "Ace of Bayes: Application of Neural Networks With Pruning", report no. 1132E, Maglegaardvej 2, DK-4000 Roskilde, Danmark

melanoma

Melanoma by type and location

Description

Data comes from a study of Malignant Melanoma involving 400 subjects.

Usage

```
data(melanoma)
```

Format

A data frame with 12 observations on the following 3 variables.

count number of cases

tumor type of tumor - a factor with levels freckle indeterminate nodular superficial

site location of tumor on the body - a factor with levels extremity head trunk

Source

Dobson A. (2002) An introduction to generalized linear models, Chapman Hall.

motorins

Third party motor insurance claims in Sweden in 1977

Description

In Sweden all motor insurance companies apply identical risk arguments to classify customers, and thus their portfolios and their claims statistics can be combined. The data were compiled by a Swedish Committee on the Analysis of Risk Premium in Motor Insurance. The Committee was asked to look into the problem of analyzing the real influence on claims of the risk arguments and to compare this structure with the actual tariff.

Usage

```
data(motorins)
```

Format

A data frame with 1797 observations on the following 8 variables.

Kilometres an ordered factor representing kilometers per year with levels 1: < 1000, 2: 1000-15000, 3: 15000-20000, 4: 20000-25000, 5: > 25000

Zone a factor representing geographical area with levels 1: Stockholm, Goteborg, Malmo with surroundings 2: Other large cities with surroundings 3: Smaller cities with surroundings in southern Sweden 4: Rural areas in southern Sweden 5: Smaller cities with surroundings in northern Sweden 6: Rural areas in northern Sweden 7: Gotland

Bonus No claims bonus. Equal to the number of years, plus one, since last claim

Make A factor representing eight different common car models. All other models are combined in class 9

Insured Number of insured in policy-years

Claims Number of claims

Payment Total value of payments in Skr

perd payment per claim

Source

<http://www.statsci.org/data/general/motorins.html>

References

Hallin, M., and Ingenbleek, J.-F. (1983). The Swedish automobile portfolio in 1977. A statistical study. *Scandinavian Actuarial Journal*, 49-64.

 neighbor

 Questionnaire study of neighborly help

Description

Subjects were asked questions in a study of neighborly help. Questions below are a subset of the full study.

Usage

`data(neighbor)`

Format

A data frame with 181 observations on the following 8 variables.

longlive About how long have you lived where you do now? Ans is a factor with levels <6mos 6-12mos 1-3yrs 3-10yrs 10yrs

wherebfr Where were you living before you moved to your present house? Ans is a factor with levels same Exeter Devon Britain Abroad

hownbly How neighborly do you think the area where you now live is? Ans is a factor with levels Vunfriendly NVfriendly Average Ffriendly Vfriendly

knowname Roughly how many people in your street, or in the streets just near you, do you know the names of? Ans is a factor with levels none 1-5 6-20 20+

callname How many of those people (not counting children) would you call by their first names? Ans is a factor with levels none 1-5 6-20 20+

age a factor with levels -18 18-30 31-50 51-65 65+

district a factor with levels 1 2 3 4

sex a factor with levels female male

Details

Exeter is a city in the county of Devon which is in Britain. The four districts can be briefly described as follows. District 1 was a long-established residential area near the city centre, with housing dating from the late nineteenth century. Originally working class, it now has a considerable middle class population with some student and other temporary accommodation. District 2 was a working-class housing estate dating from the 1930s, with mainly rented accommodation but some owner occupation. District 3 was the oldest part of a more recently developed, mainly middle-class, almost exclusively owner-occupied estate, dating from the 1960s. District 4 was the most recently developed part of a more sought-after middle-class residential area, with smaller but almost entirely owner-occupied properties dating from the 1970s and 1980s.

Source

P. Webley & S. Lea 1993, Human Relations 46, 65-76.

`nels88`*Subset of National Education Longitudinal Study 1988*

Description

Example Dataset from "Practical Regression and Anova"

Usage

```
data(nels88)
```

Format

See for yourself

Source

See Reference

References

Reference details may be found in "Practical Regression and Anova" by Julian Faraway

`nepali`*Nepali child health study*

Description

The data are a subset from public health study on Nepalese children.

Usage

```
data(nepali)
```

Format

A data frame with 1000 observations on the following 9 variables.

`id` There is a seven digit code for the child's ID: 2 digits for the panchayat number; 1 digit for the ward within panchayat; 3 digits for the household; 1 digit for child within household.

`sex` 1 = male; 2 = female

`wt` Child's weight measured in kilograms

`ht` Child's height measured in centimeters

`mage` Mother's age in years

`lit` Indicator of mother's literacy: 0 = no; 1 = yes

died The number of children the mother has had that died.
 alive The number of children the mother has ever had born alive
 age age of child

Source

West KP, Jr., LeClerq SC, Shrestha SR, Wu LS, Pradhan EK, Khatry SK, Katz J, Adhikari R, Sommer A. Effects of vitamin A on growth of vitamin A deficient children: field studies in Nepal. *J Nutr* 1997;10:1957-1965.

nes96

US 1996 national election study

Description

10 variable subset of the 1996 American National Election Study. Missing values and "don't know" responses have been listwise deleted. Respondents expressing a voting preference other than Clinton or Dole have been removed.

Usage

```
data(nes96)
```

Format

A data frame with 944 observations on the following 10 variables.

popul population of respondent's location in 1000s of people

TVnews days in the past week spent watching news on TV

selfLR Left-Right self-placement of respondent: an ordered factor with levels extremely liberal, extLib < liberal, Lib < slightly liberal, sliLib < moderate, Mod < slightly conservative, sliCon < conservative, Con < extremely conservative, extCon

ClinLR Left-Right placement of Bill Clinton (same scale as selfLR): an ordered factor with levels extLib < Lib < sliLib < Mod < sliCon < Con < extCon

DoleLR Left-Right placement of Bob Dole (same scale as selfLR): an ordered factor with levels extLib < Lib < sliLib < Mod < sliCon < Con < extCon

PID Party identification: an ordered factor with levels strong Democrat, strDem < weak Democrat, weakDem < independent Democrat, indDem < independent independentindind < indepent Republican, indRep < waek Republican, weakRep < strong Republican, strRep

age Respondent's age in years

educ Respondent's education: an ordered factor with levels 8 years or less, MS < high school dropout, HSdrop < high school diploma or GED, HS < some College, Coll < Community or junior College degree, CCdeg < BA degree, BAdeg < postgraduate degree, MAdeg

income Respondent's family income: an ordered factor with levels \$3Kminus < \$3K-\$5K < \$5K-\$7K < \$7K-\$9K < \$9K-\$10K < \$10K-\$11K < \$11K-\$12K < \$12K-\$13K < \$13K-\$14K < \$14K-\$15K < \$15K-\$17K < \$17K-\$20K < \$20K-\$22K < \$22K-\$25K < \$25K-\$30K < \$30K-\$35K < \$35K-\$40K < \$40K-\$45K < \$45K-\$50K < \$50K-\$60K < \$60K-\$75K < \$75K-\$90K < \$90K-\$105K < \$105Kplus

vote Expected vote in 1996 presidential election: a factor with levels Clinton and Dole

Source

Sapiro, Virginia, Steven J. Rosenstone, Donald R. Kinder, Warren E. Miller, and the National Election Studies. AMERICAN NATIONAL ELECTION STUDIES, 1992-1997: COMBINED FILE [Computer file]. 2nd ICPSR version. Ann Arbor, MI: University of Michigan, Center for Political Studies [producer], 1999. Ann Arbor, MI: Inter-university Consortium for Political and Social Research [distributor], 1999.

References

Found at <http://www.stat.washington.edu/quinn>

oatvar

Yields of oat varieties planted in blocks

Description

Data from an experiment to compare 8 varieties of oats. The growing area was heterogeneous and so was grouped into 5 blocks. Each variety was sown once within each block and the yield in grams per 16ft row was recorded.

Usage

`data(oatvar)`

Format

The dataset contains the following variables

yield Yield in grams per 16ft row

block Blocks I to V

variety Variety 1 to 8

Source

"Statistical Theory in Research" by R. Anderson and T. Bancroft, McGraw Hill, 1952

odor *Odor of chemical by production settings*

Description

Data from an experiment to determine the effects of column temperature, gas/liquid ratio and packing height in reducing unpleasant odor of chemical product that was being sold for household use

Usage

`data(odor)`

Format

odor Odor score
temp Temperature coded as -1, 0 and 1
gas Gas/Liquid ratio coded as -1, 0 and 1
pack Packing height coded as -1, 0 and 1

Source

"Statistical Design and Analysis of Experiments" by P. John, Macmillan, 1971

ohio *Ohio Children Wheeze Status*

Description

The ohio data frame has 2148 rows and 4 columns. The dataset is a subset of the six-city study, a longitudinal study of the health effects of air pollution.

Usage

`data(ohio)`

Format

This data frame contains the following columns:

resp an indicator of wheeze status (1=yes, 0=no)
id a numeric vector for subject id
age a numeric vector of age, 0 is 9 years old
smoke an indicator of maternal smoking at the first year of the study

References

Fitzmaurice, G.M. and Laird, N.M. (1993) A likelihood-based method for analyzing longitudinal binary responses, *Biometrika* **80**: 141–151.

orings

Spache Shuttle Challenger O-rings

Description

The 1986 crash of the space shuttle Challenger was linked to failure of O-ring seals in the rocket engines. Data was collected on the 23 previous shuttle missions. The launch temperature on the day of the crash was 31F.

Usage

`data(orings)`

Format

A data frame with 23 observations on the following 2 variables.

temp temperature at launch in degrees F

damage number of damage incidents out of 6 possible

Source

Presidential Commission on the Space Shuttle Challenger Accident, Vol. 1, 1986: 129-131.

References

S. Dalal, E. Fowlkes and B. Hoadley (1989) "Risk Analysis of the Space Shuttle: Pre-Challenger Prediction of Failure." *Journal of the American Statistical Association*. 84: 945-957.

ozone

Ozone readings in LA

Description

Example Dataset from "Practical Regression and Anova"

Usage

`data(ozone)`

Format

See for yourself

Source

See Reference

References

Reference details may be found in "Practical Regression and Anova" by Julian Faraway

parstum

Marijuana and parent alcohol and drug use

Description

445 college students were classified according to both frequency of marijuana use and parental use of alcohol and psychoactive drugs.

Usage

data(parstum)

Format

A data frame with 9 observations on the following 3 variables.

parent Number of parents using drugs or alcohol - a factor with levels Both Neither One

student Student usage of marijuana - a factor with levels Never Occasional Regular

count the number of cases

Source

Ellis, Godfrey J. and Stone, Lorene H. (1979) Marijuana Use in College: "An Evaluation of a Modeling Explanation" Youth and Society 10, 323-34

peanut

Carbon dioxide effects on peanut oil extraction

Description

The peanut data frame has 16 rows and 6 columns. Carbon dioxide effects on peanut oil extraction

Usage

```
data(peanut)
```

Format

This data frame contains the following columns:

press CO2 pressure - two levels low=0, high=1

temp CO2 temperature - two levels low=0, high=1

moist peanut moisture - two levels low=0, high=1

flow CO2 flow rate - two levels low=0, high=1

size peanut particle size - two levels low=0, high=1

solubility the amount of oil that could dissolve in the CO2

Source

Kilgo, M (1989) "An Application of Fractional Factorial Experimental Designs" *Quality Engineering*, 1, 45-54

penicillin

Penicillin yields by block and treatment

Description

Example Dataset from "Practical Regression and Anova"

Usage

```
data(penicillin)
```

Format

See for yourself

Source

See Reference

References

Reference details may be found in "Practical Regression and Anova" by Julian Faraway

pima

Diabetes survey on Pima Indians

Description

The National Institute of Diabetes and Digestive and Kidney Diseases conducted a study on 768 adult female Pima Indians living near Phoenix.

Usage

```
data(pima)
```

Format

The dataset contains the following variables

pregnant Number of times pregnant

glucose Plasma glucose concentration at 2 hours in an oral glucose tolerance test

diastolic Diastolic blood pressure (mm Hg)

triceps Triceps skin fold thickness (mm)

insulin 2-Hour serum insulin (mu U/ml)

bmi Body mass index (weight in kg/(height in metres squared))

diabetes Diabetes pedigree function

age Age (years)

test test whether the patient shows signs of diabetes (coded 0 if negative, 1 if positive)

Source

The data may be obtained from UCI Repository of machine learning databases at <http://www.ics.uci.edu/~mllearn/MLRepository.html>

pipeline	<i>NIST data on ultrasonic measurements of defects in the Alaska pipeline</i>
----------	---

Description

Researchers at National Institutes of Standards and Technology (NIST) collected data on ultrasonic measurements of the depths of defects in the Alaska pipeline in the field. The depth of the defects were then remeasured in the laboratory. These measurements were performed in six different batches. The laboratory measurements are more accurate than the in-field measurements, but more time consuming and expensive.

Usage

```
data(pipeline)
```

Format

A data frame with 107 observations on the following 3 variables.

Field measurement of depth of defect on site

Lab measurement of depth of defect in the lab

Batch the batch of measurements

Source

Office of the Director of the Institute of Materials Research (now the Materials Science and Engineering Laboratory) of NIST

pneumo	<i>Pneumoconiosis in coal miners</i>
--------	--------------------------------------

Description

The data for this example contains the number of coal miners classified by radiological examination into one of three categories of pneumonoultramicroscopicosilicovolcanoconiosis (known as pneumoconiosis for short) and by number of years spent working at the coal face divided into eight categories.

Usage

```
data(pneumo)
```

Format

A data frame with 24 observations on the following 3 variables.

freq number of miners

status pneumoconiosis status - a factor with levels mild normal severe

year number of years service (midpoint of interval)

Source

M. Aitkin and D. Anderson and B. Francis and J. Hinde (1989) "Statistical Modelling in GLIM"
Oxford University Press.

potuse

Marijuana usage by youth

Description

The National Youth Survey collected a sample of 11 to 17 year olds - 117 boys and 120 girls - asking questions about marijuana usage.

Usage

data(potuse)

Format

A data frame with 486 observations on the following 7 variables.

sex 1=Male, 2=Female

year.76 1=never used, 2=used no more than once a month, 3=used more than once a month in 1976

year.77 1=never used, 2=used no more than once a month, 3=used more than once a month in 1977

year.78 1=never used, 2=used no more than once a month, 3=used more than once a month in 1978

year.79 1=never used, 2=used no more than once a month, 3=used more than once a month in 1979

year.80 1=never used, 2=used no more than once a month, 3=used more than once a month in 1980

count Number of cases in this category

Source

ICPSR, University of Michigan

References

Lang J., McDonald, J and Smith P. (1999) "Association-Marginal Modeling of Multivariate Categorical Responses: A Maximum Likelihood Approach" JASA 94, 1161-

prostate	<i>Prostate cancer surgery</i>
----------	--------------------------------

Description

The prostate data frame has 97 rows and 9 columns. A study on 97 men with prostate cancer who were due to receive a radical prostatectomy.

Usage

```
data(prostate)
```

Format

This data frame contains the following columns:

lcvol log(cancer volume)

lweight log(prostate weight)

age age

lbph log(benign prostatic hyperplasia amount)

svi seminal vesicle invasion

lcp log(capsular penetration)

gleason Gleason score

pgg45 percentage Gleason scores 4 or 5

lpsa log(prostate specific antigen)

Source

Andrews DF and Herzberg AM (1985): Data. New York: Springer-Verlag

prplot	<i>Partial Residual Plot</i>
--------	------------------------------

Description

Makes a Partial Residual plot

Usage

```
prplot(g,i)
```

Arguments

g	An object returned from lm()
i	index of predictor

Value

none

Author(s)

Julian Faraway

Examples

```
data(stackloss)
g <- lm(stack.loss ~ .,stackloss)
prplot(g,1)
```

psid	<i>Panel Study of Income Dynamics subset</i>
------	--

Description

The Panel Study of Income Dynamics (PSID), begun in 1968, is a longitudinal study of a representative sample of U.S. individuals. The study is conducted at the Survey Research Center, Institute for Social Research, University of Michigan and is still continuing. The data represents a small subset of the total data.

Usage

```
data(psid)
```

Format

A data frame with 1661 observations on the following 6 variables.

age age in 1968

educ years of education

sex sex of individual, F or M

income annual income in dollars

year calendar year

person ID number for individual

Source

Martha S. Hill, The Panel Study of Income Dynamics: A User's Guide, Sage Publications, 1992, Newbury Park, CA.

pulp

Brightness of paper pulp depending on shift operator

Description

The pulp data frame has 20 rows and 2 columns. Data comes from an experiment to test the paper brightness depending on a shift operator.

Usage

```
data(pulp)
```

Format

This data frame contains the following columns:

bright Brightness of the pulp as measured by a reflectance meter

operator Shift operator a-d

Source

"Statistical techniques applied to production situations" F. Sheldon (1960) Industrial and Engineering Chemistry, 52, 507-509

Value

none

Author(s)

Julian Faraway

See Also

qqnorm

Examples

```
qqnorm1(rnorm(16))
```

rabbit

Rabbit weight gain by diet and litter

Description

A nutritionist studied the effects of six diets, on weight gain of domestic rabbits. From past experience with sizes of litters, it was felt that only 3 uniform rabbits could be selected from each available litter. There were ten litters available forming blocks of size three.

Usage

```
data(rabbit)
```

Format

The variables in the dataset were

treat Diet a through f

gain Weight gain

block Block (10 litters)

Source

"Experimental Design and Analysis" by M. Lentner and T. Bishop, Valley Book Company, 1986

ratdrink	<i>Rat growth weights affected by additives</i>
----------	---

Description

The data consist of 5 weekly measurements of body weight for 27 rats. The first 10 rats are on a control treatment while 7 rats have thyroxine added to their drinking water. 10 Rats have thiouracil added to their water.

Usage

```
data(ratdrink)
```

Format

A data frame with 135 observations on the following 4 variables.

wt Weight of the rat

weeks Week of the study from 0 to 4

subject the rat code number

treat treatment applied to the rat drinking water - a factor with levels control thiouracil thyroxine

Source

Unknown

rats	<i>Effect of toxic agents on rats</i>
------	---------------------------------------

Description

An experiment was conducted as part of an investigation to combat the effects of certain toxic agents.

Usage

```
data(rats)
```

Format

A data frame with 48 observations on the following 3 variables.

time survival time in tens of hours

poison the poison type - a factor with levels I II III

treat the treatment - a factor with levels A B C D

Source

Box G and Cox D (1964) "An analysis of transformations" J. Roy. Stat. Soc. Series B. **26** 211.

resceram

Shape and plate effects on current noise in resistors

Description

The resceram data frame has 12 rows and 3 columns. Shape and plate effects on current noise in resistors

Usage

```
data(resceram)
```

Format

This data frame contains the following columns:

noise current noise

shape the geometrical shape of the resistor, A, B, C or D

plate the ceramic plate on which the resistor was mounted. Only three resistors will fit on one plate.

Source

Natrella, M (1963) "Experimental Statistics" National Bureau of Standards Handbook 91, Gaithersburg MD.

salmonella

Salmonella reverse mutagenicity assay

Description

The data was collected in a salmonella reverse mutagenicity assay where the numbers of revertant colonies of TA98 Salmonella observed on each of three replicate plates for different doses of quinoline

Usage

```
data(salmonella)
```

Format

A data frame with 18 observations on the following 2 variables.

colonies numbers of revertant colonies of TA98 Salmonella

dose dose level of quinoline

Source

Breslow N.E. (1984), Extra-Poisson Variation in Log-linear Models, ApplStat, pp. 38-44.

sat

School expenditure and test scores from USA in 1994-95

Description

The sat data frame has 50 rows and 7 columns. Data were collected to study the relationship between expenditures on public education and test results.

Usage

`data(sat)`

Format

This data frame contains the following columns:

`expend` Current expenditure per pupil in average daily attendance in public elementary and secondary schools, 1994-95 (in thousands of dollars)

`ratio` Average pupil/teacher ratio in public elementary and secondary schools, Fall 1994

`salary` Estimated average annual salary of teachers in public elementary and secondary schools, 1994-95 (in thousands of dollars)

`takers` Percentage of all eligible students taking the SAT, 1994-95

`verbal` Average verbal SAT score, 1994-95

`math` Average math SAT score, 1994-95

`total` Average total score on the SAT, 1994-95

Source

"Getting What You Pay For: The Debate Over Equity in Public School Expenditures" D. Guber, Journal of Statistics Education, 1999

savings

Savings rates in 50 countries

Description

The savings data frame has 50 rows and 5 columns. The data is averaged over the period 1960-1970.

Usage

```
data(savings)
```

Format

This data frame contains the following columns:

sr savings rate - personal saving divided by disposable income

pop15 percent population under age of 15

pop75 percent population over age of 75

dpi per-capita disposable income in dollars

ddpi percent growth rate of dpi

Details

Now also appears as LifeCycleSavings in the datasets package

Source

Belsley, D., Kuh. E. and Welsch, R. (1980) "Regression Diagnostics" Wiley.

seatpos

Car seat position depending driver size

Description

Car drivers like to adjust the seat position for their own comfort. Car designers would find it helpful to know where different drivers will position the seat depending on their size and age. Researchers at the HuMoSim laboratory at the University of Michigan collected data on 38 drivers.

Usage

```
data(seatpos)
```

Format

The dataset contains the following variables

Age Age in years

Weight Weight in lbs

HtShoes Height in shoes in cm

Ht Height bare foot in cm

Seated Seated height in cm

Arm lower arm length in cm

Thigh Thigh length in cm

Leg Lower leg length in cm

hipcenter horizontal distance of the midpoint of the hips from a fixed location in the car in mm

Source

"Linear Models in R" by Julian Faraway, CRC Press, 2004

semicond

Semiconductor split-plot experiment

Description

The semicond data frame has 48 rows and 5 columns.

Format

This data frame contains the following columns:

resistance a numeric vector

ET a factor with levels 1 to 4 representing etch time.

Wafer a factor with levels 1 to 3

position a factor with levels 1 to 4

Grp an ordered factor with levels 1/1 < 1/2 < 1/3 < 2/1 < 2/2 < 2/3 < 3/1 < 3/2 < 3/3 < 4/1 < 4/2 < 4/3

Details

Also found in the SASmixed package

Source

Littel, R. C., Milliken, G. A., Stroup, W. W., and Wolfinger, R. D. (1996), *SAS System for Mixed Models*, SAS Institute (Data Set 2.2(b)).

`sexab`*Post traumatic stress disorder in abused adult females*

Description

The data for this example come from a study of the effects of childhood sexual abuse on adult females. 45 women being treated at a clinic, who reported childhood sexual abuse, were measured for post traumatic stress disorder and childhood physical abuse both on standardized scales. 31 women also being treated at the same clinic, who did not report childhood sexual abuse were also measured. The full study was more complex than reported here and so readers interested in the subject matter should refer to the original article.

Usage`data(sexab)`**Format**

The variables in the dataset are

`cpa` Childhood physical abuse on standard scale

`ptsd` Post-traumatic stress disorder on standard scale

`csa` Childhood sexual abuse - abused or not abused

Source

N. Rodriguez and S. Ryan and H. Vande Kemp and D. Foy (1997) "Postraumatic stress disorder in adult female survivors of childhood sexual abuse: A comparison study", *Journal of Consulting and Clinical Psychology*, 65, 53-59

`sexfun`*Marital sex ratings*

Description

Data from a questionnaire from 91 couples in the Tucson, Arizona area. Subjects answered the question "Sex is fun for me and my partner". The possible answers were "never or occasionally", "fairly often", "very often" and "almost always"

Usage`data(sexfun)`

Format

A data frame with 16 observations on the following 3 variables.

y the count

husband a factor with levels never fairly very always

wife a factor with levels never fairly very always

Source

Hout, M., Duncan, O. and Sobel M. (1987) Association and heterogeneity: Structural models of similarities and differences. *Sociological Methods*. 17, 145-184.

solder

Solder skips in circuit board manufacture

Description

Example Dataset from "Practical Regression and Anova"

Usage

```
data(solder)
```

Format

See for yourself

Source

See Reference

References

Reference details may be found in "Practical Regression and Anova" by Julian Faraway

 sono

Sonoluminescence

Description

The sono data frame has 16 rows and 8 columns. Sonoluminescence is the process of turning sound energy into light. An experiment was conducted to study factors affecting this process.

Usage

```
data(sono)
```

Format

This data frame contains the following columns:

Intensity Sonoluminescent light intensity

Molarity Amount of Solute. The coding is "low" for 0.10 mol and "high" for 0.33 mol.

Solute Solute type. The coding is "low" for sugar and "high" for glycerol.

pH The coding is "low" for 3 and "high" for 11.

Gas Gas type in water. The coding is "low" for helium and "high" for air.

Water Water depth. The coding is "low" for half and "high" for full.

Horn Horn depth. The coding is "low" for 5 mm and "high" for 10 mm.

Flask Flask clamping. The coding is "low" for unclamped and "high" for clamped.

Source

Eva Wilcox and Ken Inn of the NIST Physics Laboratory conducted this experiment during 1999 and published in NIST/SEMATECH e-Handbook of Statistical Methods, <http://www.itl.nist.gov/div898/handbook/>

 soybean

Germination failures for soybean seeds

Description

An experiment was conducted to compare the germination rates of the five varieties of soybean. Five plots were available.

Usage

```
data(soybean)
```

Format

A data frame with 25 observations on the following 3 variables.

variety the variety - a factor with levels arasan check fermate semesan spergon

replicate the plot - a factor with levels 1 2 3 4 5

failure the number of failures out of 100 planted seeds

Source

Snedecor G. and Cochran W. (1967) *Statistical Methods* (6th Ed) Iowa State University Press

spector

Teaching methods in Economics

Description

A study to determine the effectiveness of a new teaching method in Economics

Usage

`data(spector)`

Format

A data frame with 32 observations on the following 4 variables.

grade 1 = exam grades improved, 0 = not improved

psi 1 = student exposed to PSI (a new teach method), 0 = not exposed

tuce a measure of ability when entering the class

gpa grade point average

Source

Spector, L. and Mazzeo, M. (1980), "Probit Analysis and Economic Education", *Journal of Economic Education*, 11, 37 - 44.

speedo

Speedometer cable shrinkage

Description

Speedometer cables can be noisy because of shrinkage in the plastic casing material. An experiment was conducted to find out what caused shrinkage by screening a large number of factors. The engineers started with 15 different factors.

Usage

data(speedo)

Format

The dataset contains the following variables: (variables a-o are 2 level factors, coded "+" and "-" where "+" indicates a higher value where appropriate)

- a liner outer diameter
- b liner die
- c liner material
- d liner line speed
- e wire braid type
- f braiding tension
- g wire diameter
- h liner tension
- i liner temperature
- j coating material
- k coating die type
- l melt temperature
- m screen pack
- n cooling method
- o line speed
- y percentage shrinkage per specimen

Source

G. P. Box and S. Bisgaard and C. Fung (1988) "An explanation and critique of Taguchi's contributions to quality engineering", *Quality and reliability engineering international*, 4, 123-131

star

Star light intensities and temperatures

Description

Example Dataset from "Practical Regression and Anova"

Usage

```
data(star)
```

Format

See for yourself

Source

See Reference

References

Reference details may be found in "Practical Regression and Anova" by Julian Faraway

stat500

Scores for students in Stat500 class

Description

Example Dataset from "Practical Regression and Anova"

Usage

```
data(stat500)
```

Format

See for yourself

Source

See Reference

References

Reference details may be found in "Practical Regression and Anova" by Julian Faraway

strongx	<i>Strong interaction experiment data</i>
---------	---

Description

Example Dataset from "Practical Regression and Anova"

Usage

```
data(strongx)
```

Format

See for yourself

Source

See Reference

References

Reference details may be found in "Practical Regression and Anova" by Julian Faraway

suicide	<i>Suicide method data from the UK</i>
---------	--

Description

One year of suicide data from the United Kingdom crossclassified by sex, age and method.

Usage

```
data(suicide)
```

Format

A data frame with 36 observations on the following 4 variables.

y number of people

cause method used - a factor with levels drug (suicide by solid or liquid matter), gas, gun (guns, knives or explosives) hang (hanging, strangling, suffocating or drowning, jump other

age a factor with levels m (middle-aged) o (old) y (young)

sex a factor with levels f m

Source

Everitt B. & Dunn G. (1991) "Applied Multivariate Data Analysis" Edward Arnold

teengamb

Study of teenage gambling in Britain

Description

The teengamb data frame has 47 rows and 5 columns. A survey was conducted to study teenage gambling in Britain.

Usage

```
data(teengamb)
```

Format

This data frame contains the following columns:

sex 0=male, 1=female

status Socioeconomic status score based on parents' occupation

income in pounds per week

verbal verbal score in words out of 12 correctly defined

gamble expenditure on gambling in pounds per year

Source

Ide-Smith & Lea, 1988, Journal of Gambling Behavior, 4, 110-118

toenail

Toenail infection treatment study

Description

The data come from a Multicenter study comparing two oral treatments for toenail infection. Patients were evaluated for the degree of separation of the nail. Patients were randomized into two treatments and were followed over seven visits - four in the first year and yearly thereafter. The patients have not been treated prior to the first visit so this should be regarded as the baseline.

Usage

```
data(toenail)
```

Format

A data frame with 1908 observations on the following 5 variables.

ID ID of patient

outcome 0=none of mild separation, 1=moderate or severe

treatment the treatment A or B

month time of the visit (not exactly monthly intervals hence not round numbers)

visit the number of the visit

Source

De Backer, M., De Vroey, C., Lesaffre, E., Scheys, I., and De Keyser, P. (1998). Twelve weeks of continuous oral therapy for toenail onychomycosis caused by dermatophytes: A double-blind comparative trial of terbinafine 250 mg/day versus itraconazole 200 mg/day. *Journal of the American Academy of Dermatology*, 38, 57-63.

References

Lesaffre, E. and Spiessens, B. (2001). On the effect of the number of quadrature points in a logistic random-effects model: An example. *Journal of the Royal Statistical Society, Series C*, 50, 325-335.

G. Fitzmaurice, N. Laird and J. Ware (2004) *Applied Longitudinal Analysis*, Wiley

 troutegg

Survival of trout eggs depending on time and location

Description

Boxes of trout eggs were buried at five different stream locations and retrieved at 4 different times. The number of surviving eggs was recorded. The box was not returned to the stream.

Usage

data(troutegg)

Format

A data frame with 20 observations on the following 4 variables.

survive the number of surviving eggs

total the number of eggs in the box

location the location in the stream with levels 1 2 3 4 5

period the number of weeks after placement that the box was withdrawn levels 4 7 8 11

Source

Manly B. (1978) Regression models for proportions with extraneous variance. *Biometrie-Praximetrie*, 18, 1-18.

References

Hinde J. and Demetrio C. (1988) Overdispersion: Models and estimation. *Computational Statistics and Data Analysis*. 27, 151-170.

 truck

Truck leaf spring experiment

Description

Data on an experiment concerning the production of leaf springs for trucks. A 2^{5-1} fractional factorial experiment with 3 replicates was carried out with objective of recommending production settings to achieve a free height as close as possible to 8 inches.

Usage

`data(truck)`

Format

A data frame with 48 observations on the following 6 variables.

B furnace temperature - a factor with levels + -

C heating time - a factor with levels + -

D transfer time - a factor with levels + -

E hold-down time - a factor with levels + -

O quench oil temperature - a factor with levels + -

height leaf spring free height in inches

Source

J. J. Pignatiello and J. S. Ramberg (1985) Contribution to discussion of offline quality control, parameter design and the Taguchi method, *Journal of Quality Technology*, 17 198-206.

References

P. McCullagh and J. Nelder (1989) "Generalized Linear Models" Chapman and Hall, 2nd ed.

turtle	<i>Incubation temperature and the sex of turtles</i>
--------	--

Description

Incubation temperature and the sex of turtles

Usage

```
data(turtle)
```

Format

A data frame with 15 observations on the following 3 variables.

temp temperature in degrees centigrade

male number of male turtles hatched

female number of female turtles hatched

Details

Incubation temperature can affect the sex of turtles. There are 3 independent replicates for each temperature.

Source

Beyond Traditional Statistical Methods Copyright 2000 D. Cook, P. Dixon, W. M. Duckworth, M. S. Kaiser, K. Koehler, W. Q. Meeker and W. R. Stephenson. Developed as part of NSF/ILI grant DUE9751644.

Examples

```
data(turtle)
```

twins	<i>Twin IQs from Burt</i>
-------	---------------------------

Description

Example Dataset from "Practical Regression and Anova"

Usage

```
data(twins)
```

Format

See for yourself

Source

See Reference

References

Reference details may be found in "Practical Regression and Anova" by Julian Faraway

uncviet

UNC student opinions about the Vietnam War

Description

A student newspaper conducted a survey of student opinions about the Vietnam War in May 1967. Responses were classified by sex, year in the program and one of four opinions. The survey was voluntary.

Usage

```
data(uncviet)
```

Format

A data frame with 40 observations on the following 4 variables.

y the count

policy a factor with levels A (defeat power of North Vietnam by widespread bombing and land invasion) B (follow the present policy) C (withdraw troops to strong points and open negotiations on elections involving the Viet Cong) D (immediate withdrawal of all U.S. troops)

sex a factor with levels Female Male

year a factor with levels Fresh Grad Junior Senior Soph

Source

M. Aitkin and D. Anderson and B. Francis and J. Hinde (1989) "Statistical Modelling in GLIM" Oxford University Press.

uswages

Weekly wages of US male workers in 1988

Description

The uswages data frame has 2000 rows and 10 columns. Weekly Wages for US male workers sampled from the Current Population Survey in 1988.

Usage

```
data(uswages)
```

Format

This data frame contains the following columns:

wage Real weekly wages in dollars (deflated by personal consumption expenditures - 1992 base year)

educ Years of education

exper Years of experience

race 1 if Black, 0 if White (other races not in sample)

smsa 1 if living in Standard Metropolitan Statistical Area, 0 if not

ne 1 if living in the North East

mw 1 if living in the Midwest

we 1 if living in the West

so 1 if living in the South

pt 1 if working part time, 0 if not

Source

Bierens, H.J., and D. Ginther (2001): "Integrated Conditional Moment Testing of Quantile Regression Models", *Empirical Economics* 26, 307-324

vif

vif

Description

Computes the variance inflation factors

Usage

```
vif(object)
```

Arguments

object a data matrix (design matrix without intercept) or a model object

Value

variance inflation factors

Author(s)

Julian Faraway

Examples

```
data(stackloss)
vif(stackloss[, -4])
# Air.Flow Water.Temp Acid.Conc.
# 2.9065 2.5726 1.3336
```

vision

Vision acuity tests

Description

Example Dataset from "Practical Regression and Anova"

Usage

```
data(vision)
```

Format

See for yourself

Source

See Reference

References

Reference details may be found in "Practical Regression and Anova" by Julian Faraway

wafer	<i>resistivity of wafer in semiconductor experiment</i>
-------	---

Description

A full factorial experiment with four two-level predictors.

Usage

```
data(wafer)
```

Format

A data frame with 16 observations on the following 5 variables.

x1 a factor with levels - +

x2 a factor with levels - +

x3 a factor with levels - +

x4 a factor with levels - +

resist Resistivity of the wafer

Source

Myers, R. and Montgomery D. (1997) A tutorial on generalized linear models, *Journal of Quality Technology*, 29, 274-291.

wavesolder	<i>Defects in a wave soldering process</i>
------------	--

Description

Components are attached to an electronic circuit card assembly by a wave-soldering process. The soldering process involves baking and preheating the circuit card and then passing it through a solder wave by conveyor. Defect arise during the process. Design is 2^{7-3} with 3 replicates.

Usage

```
data(wavesolder)
```

Format

A data frame with 16 observations on the following 10 variables.

y1 Number of defects in the first replicate

y2 Number of defects in the second replicate

y3 Number of defects in the third replicate

prebake prebake condition - a factor with levels 1 2

flux flux density - a factor with levels 1 2

speed conveyor speed - a factor with levels 1 2

preheat preheat condition - a factor with levels 1 2

cooling cooling time - a factor with levels 1 2

agitator ultrasonic solder agitator - a factor with levels 1 2

temp solder temperature - facctor with levels 1 2

Source

L. Condra (1993) Reliability improvement with design of experiments. Marcel Dekker, NY.

References

M. Hamada and J. Nelder (1997) Generalized linear models for quality improvement experiments, Journal of Quality Technology, 29, 292-304

wbca

Wisconsin breast cancer database

Description

Data come from a study of breast cancer in Wisconsin. There are 681 cases of potentially cancerous tumors of which 238 are actually malignant. Determining whether a tumor is really malignant is traditionally determined by an invasive surgical procedure. The purpose of this study was to determine whether a new procedure called fine needle aspiration which draws only a small sample of tissue could be effective in determining tumor status.

Usage

`data(wbca)`

Format

A data frame with 681 observations on the following 10 variables.

Class 0 if malignant, 1 if benign

Adhes marginal adhesion

BNucl bare nuclei

Chrom bland chromatin

Epith epithelial cell size

Mitos mitoses

NNucl normal nucleoli

Thick clump thickness

UShap cell shape uniformity

USize cell size uniformity

Details

The predictor values are determined by a doctor observing the cells and rating them on a scale from 1 (normal) to 10 (most abnormal) with respect to the particular characteristic.

Source

Bennett, K.,P., and Mangasarian, O.L., Neural network training via linear programming. In P. M. Pardalos, editor, *Advances in Optimization and Parallel Computing*, pages 56-57. Elsevier Science, 1992

weldstrength	<i>welding strength DOE</i>
--------------	-----------------------------

Description

An experiment to investigate factors affecting welding strength.

Usage

```
data(weldstrength)
```

Format

A data frame with 16 observations on the following 10 variables.

Rods a 0-1 predictor

Drying a 0-1 predictor

Material a 0-1 predictor

Thickness a 0-1 predictor

Angle a 0-1 predictor
Opening a 0-1 predictor
Current a 0-1 predictor
Method a 0-1 predictor
Preheating a 0-1 predictor
Strength The welding strength

Source

G. Box and R. Meyer (1986) Dispersion effects from fractional designs, *Technometrics*, 28, 19-27

wheat

Insect damage to wheat varieties

Description

Example Dataset from "Practical Regression and Anova"

Usage

```
data(wheat)
```

Format

See for yourself

Source

See Reference

References

Reference details may be found in "Practical Regression and Anova" by Julian Faraway

Index

*Topic **datasets**

aatemp, 4
abrasion, 5
aflatoxin, 5
africa, 6
alfalfa, 7
amlxray, 7
babyfood, 8
beetle, 9
bliss, 9
breaking, 10
broccoli, 10
cathedral, 11
chicago, 12
chiczip, 12
chmiss, 13
choccake, 13
chredlin, 14
clot, 15
cmob, 15
cns, 16
coagulation, 17
composite, 17
cornnit, 18
corrosion, 18
cpd, 19
ctsib, 20
death, 20
debt, 21
diabetes, 22
dicentric, 23
divusa, 24
drugpsy, 24
dvisits, 25
eco, 26
eggprod, 27
eggs, 27
epilepsy, 28
esdcomp, 28
exa, 29
exb, 29
eyegrade, 30
fat, 30
femsmoke, 31
fpe, 32
fruitfly, 33
gala, 33
gavote, 34
haireye, 35
happy, 36
hormone, 37
hprice, 37
hsb, 38
infmort, 39
irrigation, 40
jsp, 41
kanga, 41
lawn, 42
leafblotch, 42
mammalsleep, 44
meatspec, 45
melanoma, 46
motorins, 47
neighbor, 48
nels88, 49
nepali, 49
nes96, 50
oatvar, 51
odor, 52
ohio, 52
orings, 53
ozone, 53
parstum, 54
peanut, 55
penicillin, 55
pima, 56
pipeline, 57
pneumo, 57

- potuse, 58
- prostate, 59
- psid, 60
- pulp, 61
- pvc, 62
- rabbit, 63
- ratdrink, 64
- rats, 64
- resceram, 65
- salmonella, 65
- sat, 66
- savings, 67
- seatpos, 67
- semicond, 68
- sexab, 69
- sexfun, 69
- solder, 70
- sono, 71
- soybean, 71
- spector, 72
- speedo, 73
- star, 74
- stat500, 74
- strongx, 75
- suicide, 75
- teengamb, 76
- toenail, 76
- troutegg, 77
- truck, 78
- turtle, 79
- twins, 79
- uncviet, 80
- uswages, 81
- vision, 82
- wafer, 83
- wavesolder, 83
- wbca, 84
- weldstrength, 85
- wheat, 86
- *Topic **regression**
 - Cpplot, 19
 - maxadjr, 45
 - prplot, 60
- aatemp, 4
- abrasion, 5
- aflatoxin, 5
- africa, 6
- alfalfa, 7
- amlxray, 7
- babyfood, 8
- beetle, 9
- bliss, 9
- breaking, 10
- broccoli, 10
- cathedral, 11
- chicago, 12
- chiczip, 12
- chmiss, 13
- choccake, 13
- chredlin, 14
- clot, 15
- cmob, 15
- cns, 16
- coagulation, 17
- composite, 17
- cornnit, 18
- corrosion, 18
- cpd, 19
- Cpplot, 19
- ctsib, 20
- death, 20
- debt, 21
- diabetes, 22
- dicentric, 23
- divusa, 24
- drugpsy, 24
- dvisits, 25
- eco, 26
- eggprod, 27
- eggs, 27
- epilepsy, 28
- esdcomp, 28
- exa, 29
- exb, 29
- eyegrade, 30
- fat, 30
- femsmoke, 31
- fpe, 32
- fruitfly, 33
- gala, 33
- gavote, 34

haireye, 35
halfnorm, 35
happy, 36
hormone, 37
hprice, 37
hsb, 38

ilogit, 39
infmort, 39
irrigation, 40

jsp, 41

kanga, 41

lawn, 42
leafblotch, 42
logit, 43

mammalsleep, 44
maxadjr, 45
meatspec, 45
melanoma, 46
motorins, 47

neighbor, 48
nels88, 49
nepali, 49
nes96, 50

oatvar, 51
odor, 52
ohio, 52
orings, 53
ozone, 53

parstum, 54
peanut, 55
penicillin, 55
pima, 56
pipeline, 57
pneumo, 57
potuse, 58
prostate, 59
prplot, 60
psid, 60
pulp, 61
pvc, 62

qqnorm1, 62

rabbit, 63
ratdrink, 64
rats, 64
resceram, 65

salmonella, 65
sat, 66
savings, 67
seatpos, 67
semicond, 68
sexab, 69
sexfun, 69
solder, 70
sono, 71
soybean, 71
specter, 72
speedo, 73
star, 74
stat500, 74
strongx, 75
suicide, 75

teengamb, 76
toenail, 76
troutegg, 77
truck, 78
turtle, 79
twins, 79

uncviet, 80
uswages, 81

vif, 81
vision, 82

wafer, 83
wavesolder, 83
wbca, 84
weldstrength, 85
wheat, 86